

Superconvergence Relations ¹

Reinhard Oehme ²

*Enrico Fermi Institute and Department of Physics
University of Chicago
Chicago, Illinois, 60637, USA*

Abstract

For a limited number of matter fields, the discontinuity of the transverse gauge field propagator can satisfy an exact sum rule. With controlled and limited gauge dependence, this superconvergence relation is of physical interest.

¹For the ‘*Concise Encyclopedia of SUPERSYMMETRY*’,
Kluwer Academic Publishers, Dordrecht, (Editors: Jon Bagger, Steven Duplij and Warren Siegel)
2001.

²E-mail: oehme@theory.uchicago.edu

SUPERCONVERGENCE RELATIONS are sum rules for the discontinuity $\rho(k^2)$ of a structure function $D(k^2)$ in gauge theories [1]. The transverse gauge field propagator is considered here as a characteristic example. From Lorentz covariance and a minimal spectral condition, it follows that $D(k^2)$ is the boundary value of a function which is holomorphic in the complex k^2 plane with a cut along the positive real axis. A priori, with the space-time propagator being a tempered distribution, this function is bounded by a polynomial for $k^2 \rightarrow \infty$. But for regions in the parameter space where the theory is asymptotically free, the asymptotic forms can be calculated in terms of the weak coupling limit. Using renormalization group methods, one finds for the structure function

$$-k^2 D(k^2, \kappa^2, g, \alpha) \simeq \frac{\alpha}{\alpha_0} + C(g^2, \alpha) \left(-\beta_0 \ln \frac{k^2}{\kappa^2} \right)^{-\gamma_{00}/\beta_0} + \dots, \quad (1)$$

and the asymptotic term for the discontinuity along the positive, real k^2 -axis is given by

$$-k^2 \rho(k^2, \kappa^2, g, \alpha) \simeq \frac{\gamma_{00}}{\beta_0} C(g^2, \alpha) \left(-\beta_0 \ln \frac{k^2}{|\kappa^2|} \right)^{-\gamma_{00}/\beta_0 - 1} + \dots. \quad (2)$$

Here $\kappa^2 < 0$ is the normalization point, and

$$\gamma(g^2, \alpha) = (\gamma_{00} + \alpha \gamma_{01}) g^2 + \dots, \quad (3)$$

$$\beta(g^2) = \beta_0 g^4 + \dots \quad (4)$$

are the limits $g^2 \rightarrow 0$ of the anomalous dimension and the renormalization group function, while $\alpha_0 = -\gamma_{00}/\gamma_{01}$. It is important that the functional form of the asymptotic discontinuity is independent of the gauge parameter $\alpha \geq 0$.

Given the analytic and asymptotic properties of $D(k^2)$, this function satisfies the dispersion relation

$$D(k^2) = \int_{-0}^{\infty} dk'^2 \frac{\rho(k'^2)}{(k'^2 - k^2)}. \quad (5)$$

For the special case $\gamma_{00}/\beta_0 > 0$, there is sufficient boundedness for the validity of the Superconvergence Relation [1]

$$\int_{-0}^{\infty} dk^2 \rho(k^2, \kappa^2, g, \alpha) = \frac{\alpha}{\alpha_0} , \quad (6)$$

which is most directly useful in the Landau gauge, where $\alpha = 0$:

$$\int_{-0}^{\infty} dk^2 \rho(k^2, \kappa^2, g, 0) = 0 . \quad (7)$$

The superconvergence relations are exact, and are not valid order by order in perturbation theory. They depend upon short- and long-distance properties of the theory. Given $\beta_0 < 0$, it is the sign of the anomalous dimension coefficient γ_{00} which determines the existence or non-existence of superconvergence in all gauges. In connection with the BRST cohomology, the superconvergence relations can be used in order to argue for the absence of the transverse gauge field quanta from the physical state space of the theory (confinement) [2]. Hence $\gamma_{00} = 0$ indicates a phase transition similar to $\beta_0 = 0$.

For SQCD with the gauge group $SU(N_C)$ and N_F flavors, the zero of $\gamma_{00}(N_F, N_C)$ is at $N_F = \frac{3}{2}N_C$, and that of $\beta_0(N_F, N_C)$ at $N_F = 3N_C$. These values of N_F define the lower and the upper limit of the conformal window [3]. The superconvergence relations are valid for $N_F < \frac{3}{2}N_C$. As described above, for these values of N_F , the gauge quanta are not in the physical state space.

There are also superconvergence relations for non-supersymmetric theories, and results similar to those for SQCD were obtained earlier for QCD [2]. With the same gauge group $SU(N_C)$ and N_F flavors, one has superconvergence and confinement for $N_F < \frac{13}{4}N_C$. The window is given by $\frac{13}{4}N_C < N_F < \frac{22}{4}N_C$.

It is of interest to consider the possibility of superconvergence relations for theories with space-time non-commutativity. When time is involved, the integrand of a possible sum rule would be expected to have contributions from discontinuities of tachyonic branch cuts which are not directly associated with the input field content of the theory [4]. Dispersion relations for scattering amplitudes are also modified by additional singularities associated with violations of locality [5].

The new singularities are very different from the anomalous thresholds, which can be present even in local field theories. These thresholds are well understood as structure singularities associated with the composite structure of the particles involved in the theory [6]. If followed into secondary Riemann sheets by changing the mass variables,

the anomalous thresholds are seen to be ordinary crossed-channel thresholds (left hand cuts) of amplitudes connected by unitarity to the one under consideration. There are no structure singularities in the gauge field propagator. With appropriately generalized BRST-methods, possible superconvergence relations could be of interest for the interpretation of non-commutative gauge theories.

BIBLIOGRAPHY.

- [1] R. Oehme, W. Zimmermann, Phys. Rev. **D21** (1980) 471; Phys. Rev. **D21** (1980) 1661; R. Oehme, Phys. Lett. **B252** (1990) 641; R. Oehme, W. Xu Phys. Lett. **B333** (1994) 172, hep-th/9406081; **B384** (1994) 269, hep-th/9604021.
- [2] R. Oehme, Phys. Rev. **D42** (1990) 4209; Phys. Lett. **B195** (1987) 60; K. Nishijima, Prog.Theor.Phys. **75** (1986) 1221.
- [3] R. Oehme, ‘Superconvergence, Supersymmetry and Conformal Invariance’, in Leite Lopes Festschrift, pp. 443-457, World Scientific 1988 (KEK Library Scan, HEP SPIRES SLAC: Oehme, R.); N. Seiberg, Nucl. Phys. **B435** (1995) 129, hep-th/9411149; R. Oehme, Phys. Lett. **B399** (1997) 67, hep-th/9701012; ‘Superconvergence and Duality’, Concise Encyclopedia of SUPERSYMMETRY, Kluwer 2001, hep-th 0101021.
- [4] J. Gomis and T. Mehen, Nucl. Phys. **B591** (2000) 265, hep-th/0005129; L. Alvarez-Gaume, J. Barbón and R. Zwicky, hep-th/0103069; (these papers contain further references).
- [5] R. Oehme, ‘Forward Dispersion Relations and Microcausality’, in Quanta, Wentzel Festschrift (University of Chicago Press, 1970) pp. 309-337; Phys. Rev. **100** (1955) 1503; (these papers contain further references).
- [6] R. Oehme, ‘The Compound Structure of Elementary Particles’, in Werner Heisenberg und die Physik unserer Zeit, Festschrift (Verlag Friedrich Vieweg und Sohn, Braunschweig, 1961) pp. 240-259; Phys. Rev. **111** (1958) 1430; Phys. Rev. **121** (1961) 1840; Nuovo Cimento **13** (1959) 778; (these papers contain further references).